arXiv:1609.05493v1 [math.CO] 18 Sep 2016

RATIONALITY IN MAP AND HYPERMAP ENUMERATION BY GENUS

M. KAZARIAN, P. ZOGRAF

Abstract. Generating functions for a fixed genus map and hypermap enumeration become rational after a simple explicit change of variables. Their numerators are polynomials with integer coefficients that obey a differential recursion, and denominators are products of powers of explicit linear functions.

1. Introduction

By a *map* or a *ribbon graph* we understand a finite connected graph with prescribed cyclic orders of half-edges at each vertex. It also can be realized as the 1-skeleton of a polygonal partition of a closed orientable surface. The genus g of a map (ribbon graph) satisfies the Euler formula

$$2-2g = \#v - \#e + \#f\,,$$

where $\#v, \#e, \#f$ are the numbers of vertices, edges and faces of the map respectively. By a *hypermap* we understand a bicolored map, i. e. a map whose faces are properly colored in two colors (say, white and black) so that no adjacent faces have the same color. The dual graph to a hypermap is a bipartite ribbon graph, or a *Grothendieck's "dessin d'enfant"* [1].

We are interested in the weighted count of maps and hypermaps, where the weights are reciprocal to the orders of the corresponding automorphism groups. This is equivalent to counting *rooted* maps and hypermaps (i. e. those with a marked half-edge). The passage from the rooted count to the unrooted one is known, cf. [8], [9].

Denote by $\tilde{c}_{g,n}$ (resp. $c_{g,n}$) the number of rooted maps (resp. hypermaps) of genus g with n edges (darts), and consider the genus g generating functions

$$\widetilde{C}_g(s) = \sum_{n=2g}^{\infty} \tilde{c}_{g,n} s^n \quad g \geq 0\,, \tag{1}$$

$$C_g(s) = \sum_{n=2g+1}^{\infty} c_{g,n} s^n \quad g \geq 0\,. \tag{2}$$

The classical problem that goes back to Tutte [10] (or even earlier) is to compute the numbers $c_{g,n}$ and $\tilde{c}_{g,n}$. Effective algorithms for computing these numbers first appeared in [12] (for maps) and in [13] (for hypermaps)[2]. Recursions for the numbers $c_{g,n}$ and $\tilde{c}_{g,n}$, and differential equations for the generating functions $C_g(s)$ and

2010 *Mathematics Subject Classification.* 05C30, 37K10.

Key words and phrases. enumeration of maps (ribbon graphs), enumeration of hypermaps (Grothendieck's "dessins d'enfants").

[1] As observed by Grothendieck, the absolute Galois group $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ naturally acts on dessins (hypermaps); we refer the reader to [7] for details.

[2] The effective enumeration of 1-vertex maps was obtained in [4], and of 1-vertex hypermaps – in [5] and, independently, in [1]. Enumeration of 1-vertex maps (or genus g gluings of a $2n$-gon) was a crucial point in computing the Euler characteristic of the moduli space of algebraic curves in [4].

$\widetilde{C}_g(s)$ were first obtained in [6] (cf. also [2] for an alternative approach to map enumeration).

In this note we show that the generating functions $C_g(s)$ and $\widetilde{C}_g(s)$ become rational functions after simple explicit changes of variable s. Their numerators are then polynomials with integer coefficients that obey a differential recursion, and denominators are products of powers of explicit linear functions.

2. Main results

We start with the case of hypermaps (Grothendieck's dessins d'enfants).

Theorem 1. *Under the substitution $s = t(1-2t)$ we have*

$$C_0(t(1-2t)) = \frac{t(1-3t)}{(1-2t)^2}\,,$$
$$C_1(t(1-2t)) = \frac{t^3}{(1-t)(1-4t)^2}\,,$$
$$C_g(t(1-2t)) = \frac{P_g(t)}{(1-t)^{4g-3}(1-4t)^{5g-3}}\,, \quad g \geq 2, \tag{3}$$

where $P_g(t) = \sum_{i=2g+1}^{9g-7} p_{g,i}\, t^i$ is a polynomial with integer coefficients and $p_{g,2g+1} = \frac{(2g)!}{g+1}$. The polynomials $P_g(t)$ can be computed recursively by Eq. (9).

Remark 1. The polynomials $P_g(t)$ for $g = 2, 3$ are:

$$P_2(t) = 8t^5 - 92t^6 + 464t^7 - 1316t^8 + 2204t^9 - 2048t^{10} + 816t^{11},$$
$$\begin{aligned} P_3(t) = {} & 180t^7 - 3648t^8 + 35424t^9 - 218944t^{10} + 958160t^{11} - 3102528t^{12} \\ & + 7503664t^{13} - 13310768t^{14} + 16365216t^{15} - 11823680t^{16} + 117916t^{17} \\ & + 6614784t^{18} - 6008320t^{19} + 1823744t^{20}. \end{aligned}$$

In principle, they can be computed for much larger values of g.

Remark 2. Similar result was independently obtained in [3] by a different (more complicated) method.

Proof. To prove the theorem, we recall a specialization of the Kadomtsev-Petviashvili (KP) equation for the hypermap count derived in [6]:

$$\begin{aligned} (sC_g)' = {} & 3(2s^2C_g' + sC_g) + 3s^3C_g' + s^3(s(sC_{g-1})')'' \\ & + s^3 \sum_{i=0}^{g} (4C_i + 6sC_i')C_{g-i}' + 2s\delta_{g,0}\,, \end{aligned} \tag{4}$$

where the prime $'$ stands for the derivative $\frac{d}{ds}$. This equation is just the differential form of the recursion (11) in [6] for $t = u = v = 1$:

$$\begin{aligned} (n+1)c_{g,n} = {} & 3(2n-1)\,c_{g,n-1} + (n-2)\,c_{g,n-2} + (n-1)^2(n-2)\,c_{g-1,n-2} \\ & + \sum_{i=0}^{g} \sum_{j=1}^{n-3} (4+6j)(n-2-j)\,c_{i,j}\,c_{g-i,n-2-j}\,. \end{aligned} \tag{5}$$

For $g \geq 1$ we can further rewrite Eq. (4) as the differential recursion

$$\begin{aligned} & (s - 6s^2 - 3s^3 - 4s^3C_0 - 12s^4C_0')\,C_g' + (1 - 3s - 4s^3C_0')C_g \\ & = s^5C_{g-1}''' + 5s^4C_{g-1}'' + 4s^3C_{g-1}' + s^3 \sum_{i=1}^{g-1} (4C_i + 6sC_i')C_{g-i}'\,. \end{aligned} \tag{6}$$

For $g = 0$ we get an ordinary differential equation that can be solved explicitly:

$$C_0(s) = \frac{-1 + 12s - 24s^2 + (1-8s)^{3/2}}{32s^2}\,.$$

It is easy to see that the substitution $s = t(1-2t)$ considerably simplifies C_0 and makes it a rational function, namely

$$C_0(t(1-2t)) = \frac{t(1-3t)}{(1-2t)^2}$$

(cf. [11]). Substituting $s = t(1-2t)$ in (6), we get

$$\begin{aligned} &t(1-t)^2(1-2t)\,\dot{C}_g + (1-t)(1-2t+4t^2)\,C_g \\ &= t^3(1-2t)^3\left(D_t C_{g-1} + \frac{1}{(1-4t)^2}\sum_{i=1}^{g-1}\left(4(1-4t)C_i + 6t(1-2t)\dot{C}_i\right)\dot{C}_{g-i}\right), \end{aligned} \tag{7}$$

where $\dot{C}_g = \frac{d}{dt}C_g(t(1-2t))$ and

$$\begin{aligned} D_t = \frac{t^2(1-2t)^2}{(1-4t)^3}\cdot\frac{d^3}{dt^3} + \frac{t(1-2t)(5-28t+56t^2)}{(1-4t)^4}\cdot\frac{d^2}{dt^2} \\ + \frac{4(1-11t+58t^2-144t^3+144t^4)}{(1-4t)^5}\cdot\frac{d}{dt}\,. \end{aligned} \tag{8}$$

Assuming that $C_0, \ldots, C_{g-1}$ are known, we can think of (7) as an ODE for C_g. The integrating factor for this equation is $\frac{1-t}{(1-2t)^3}$, so that we get from (7)

$$\begin{aligned} &\frac{d}{dt}\left(\frac{t(1-t)^3}{(1-2t)^2}\,C_g\right) \\ &\quad = t^3(1-t)\left(D_t C_{g-1} + \frac{1}{(1-4t)^2}\sum_{i=1}^{g-1}\left(4(1-4t)C_i + 6t(1-2t)\dot{C}_i\right)\dot{C}_{g-i}\right), \end{aligned}$$

or, equivalently,

$$\begin{aligned} &C_g(t(1-2t) = \frac{(1-2t)^2}{t(1-t)^3} \\ &\times \int t^3(1-t)\left(D_t C_{g-1} + \frac{1}{(1-4t)^2}\sum_{i=1}^{g-1}\left(4(1-4t)C_i + 6t(1-2t)\dot{C}_i\right)\dot{C}_{g-i}\right)dt\,. \end{aligned} \tag{9}$$

Since by definition $C_g(0) = 0$ for all $g \geq 0$, Eq. (9) determines C_g uniquely in terms of $C_0, \ldots, C_{g-1}$. In particular, this equation immediately yields

$$\begin{aligned} C_1(t(1-2t)) &= \frac{t^3}{(1-t)(1-4t)^2}\,, \\ C_2(t(1-2t)) &= \frac{8t^5 - 92t^6 + 464t^7 - 1316t^8 + 2204t^9 - 2048t^{10} + 816t^{11}}{(1-t)^5(1-4t)^7}\,. \end{aligned}$$

Let us show that $C_g(t(1-2t))$ has the form (3) for any $g \geq 3$. We will use the elementary formula

$$\frac{d}{dt}\left(\frac{t^\alpha}{(1-t)^\beta(1-4t)^\gamma}\right) = \frac{\alpha t^{\alpha-1} + (-5\alpha+\beta+4\gamma)t^\alpha + 4(\alpha-\beta-\gamma)t^{\alpha+1}}{(1-t)^{\beta+1}(1-4t)^{\gamma+1}}\,. \tag{10}$$

Then we have

$$D_t C_{g-1} = \frac{(2g-1)(2g)^2 p_{g-1,2g-1}t^{2g-2} + \ldots - 256 p_{g-1,9g-16}t^{9g-9}}{(1-t)^{4g-4}(1-4t)^{5g-2}} \tag{11}$$

and

$$\frac{1}{(1-4t)^2}\sum_{i=1}^{g-1}\Big(4(1-4t)C_i+6t(1-2t)\dot{C}_i\Big)\,\dot{C}_{g-i} \\ =\frac{r_g t^{2g+1}+\ldots+256p_{g-1,9g-16}t^{9g-9}}{(1-t)^{4g-4}(1-4t)^{5g-2}}\,, \tag{12}$$

where r_g is some constant. Notice that the top degree term in the numerator on the right hand side of (12) comes entirely from the product $C_1\dot{C}_{g-1}$. Multiplying both sides of (11) and (12) by $t^3(1-t)$ and taking their sum we get that the integrand in (9) has the form

$$\frac{Q_g(t)}{(1-t)^{4g-5}(1-4t)^{5g-2}}\,, \tag{13}$$

where $Q_g(t)=\sum_{i=2g+1}^{9g-7} q_{g,i}t^i$ is a polynomial with $q_{g,2g+1}=(2g-1)(2g)^2p_{g-1,2g-1}$. Therefore, we can rewrite (9) as

$$C_g(t(1-2t))=\frac{(1-2t)^2}{t(1-t)^3}\int\frac{Q_g(t)}{(1-t)^{4g-5}(1-4t)^{5g-2}}dt\,. \tag{14}$$

To perform integration in (14) we decompose the integrand into the sum

$$\frac{Q_g(t)}{(1-t)^{4g-5}(1-4t)^{5g-2}}=a+\sum_{i=2}^{4g-5}\frac{a_i}{(1-t)^i}+\sum_{j=2}^{5g-2}\frac{b_j}{(1-4t)^j}\,. \tag{15}$$

Note that no terms of the form $\frac{a_1}{1-t}$ or $\frac{b_1}{1-4t}$ can appear in the right hand side of (15) because the Taylor series expansion of the left hand side of (14) has integer coefficients[3]. Integrating we obtain

$$\int\frac{Q_g(t)}{(1-t)^{4g-5}(1-4t)^{5g-2}}dt=at+b+\sum_{i=1}^{4g-6}\frac{a_{i+1}}{i}\cdot\frac{1}{(1-t)^i}+\sum_{j=1}^{5g-3}\frac{b_{j+1}}{4j}\cdot\frac{1}{(1-4t)^j}\,, \tag{16}$$

where the condition $C_g(0)=0$ implies

$$b=-\sum_{i=1}^{4g-6}\frac{a_i}{i}-\sum_{j=2}^{5g-3}\frac{b_{j+1}}{4j}\,.$$

Multiplying the right hand side of (16) by $(1-t)^{4g-6}(1-4t)^{5g-3}$ we get a polynomial of the form $R_g(t)=\sum_{i=2g+2}^{9g-8} r_{g,i}t^i$. To complete the proof we put $P_g(t)=\frac{(1-2t)^2}{t}R_g(t)$ and notice that $p_{g,2g+1}=\frac{(2g-1)(2g)^2}{2g+2}\,p_{g-1,2g-1}$. Moreover, we see that $t=1/2$ is a root of $P_g(t)$ of multiplicity 2 provided $g\geq 2$.[4] □

Now we continue with map enumeration.

Theorem 2. *Under the substitution $s=t(1-3t)$ we have*

$$\widetilde{C}_0(t(1-3t))=\frac{1-4t}{(1-3t)^2}\,,$$

$$\widetilde{C}_1(t(1-3t))=\frac{t^2}{(1-2t)(1-6t)^2}\,,$$

[3]We owe this observation to F. Petrov.

[4]Numerically, we also have $p_{g,9g-7}\neq 0$, $P_g(1)\neq 0$, $P_g(1/4)\neq 0$. In principle, this can be verified along the same lines as above, but computations become too cumbersome to reproduce them here.

$$\widetilde{C}_g(t(1-3t)) = \frac{\widetilde{P}_g(t)}{(1-2t)^{3g-2}(1-6t)^{5g-3}}\,, \quad g \geq 2, \tag{17}$$

where $\widetilde{P}_g(t) = \sum_{i=2g}^{8g-6} \tilde{p}_{g,i}\, t^i$ *with* $\tilde{p}_{g,2g} = \frac{(4g-1)!!}{2g+1}$. *The polynomials* $\widetilde{P}_g(t)$ *can be computed recursively by Eq. (23).*

Remark 3. The polynomials $\widetilde{P}_g(t)$ for $g = 2, 3$ are:

$$\begin{aligned}
\widetilde{P}_2(t) &= 21t^4 - 336t^5 + 2334t^6 - 9108t^7 + 21177t^8 - 27756t^9 + 15876t^{10},\\
\widetilde{P}_3(t) &= 1485t^6 - 41184t^7 + 539073t^8 - 4483458t^9 + 26893989t^{10} - 124232004t^{11}\\
&\quad + 453861279t^{12} - 1307353122t^{13} + 2897271774t^{14} - 4737605112t^{15}\\
&\quad + 5355443952t^{16} - 3723895296t^{17} + 1197496224t^{18}.
\end{aligned}$$

Like in the case of hypermaps, they can be computed for much larger values of g.

Proof. The proof of Theorem 2 is very similar to that of Theorem 1, so we will outline only its main steps. We recall a specialization of the Kadomtsev-Petviashvili (KP) equation for the map count derived in [6]:

$$\begin{aligned}
(s\widetilde{C}_g)' &= 4(2s^2\widetilde{C}_g' + s\widetilde{C}_g) + 2s^3(2s(s\widetilde{C}_{g-1})' + s\widetilde{C})'' + s^2(2s(s\widetilde{C}_{g-1})' + s\widetilde{C})'\\
&\quad + 3s^2\sum_{i=0}^{g}(\widetilde{C}_i + 2s\widetilde{C}_i')(\widetilde{C}_{g-i} + 2s\widetilde{C}_{g-i}') + \delta_{g,0}\,,
\end{aligned} \tag{18}$$

where the prime $'$ stands for the derivative $\frac{d}{ds}$. This equation is just a differential form of the recursion (16) in [6] for $t = u = 1$:

$$\begin{aligned}
(n+1)\tilde{c}_{g,n} &= 4(2n-1)\,\tilde{c}_{g,n-1} + (2n-1)(2n-3)(n-1)\,\tilde{c}_{g-1,n-2}\\
&\quad + 3\sum_{i=0}^{g}\sum_{j=0}^{n-2}(2j+1)(2(n-2-j)+1)\,\tilde{c}_{i,j}\,\tilde{c}_{g-i,n-2-j}\,.
\end{aligned} \tag{19}$$

For $g \geq 1$ we can further rewrite Eq. (18) as the differential recursion

$$\begin{aligned}
&(s - 8s^2 - 12s^3\widetilde{C}_0 - 24s^4\widetilde{C}_0')\,\widetilde{C}_g' + (1 - 4s - 6s^2\widetilde{C}_0 - 12s^3\widetilde{C}_0')\,\widetilde{C}_g\\
&= 4s^5\widetilde{C}_{g-1}''' + 24s^4\widetilde{C}_{g-1}'' + 27s^3\widetilde{C}_{g-1}' + 3s^2\widetilde{C}_{g-1} + 3s^2\sum_{i=0}^{g}(\widetilde{C}_i + 2s\widetilde{C}_i')(\widetilde{C}_{g-i} + 2s\widetilde{C}_{g-i}')\,.
\end{aligned} \tag{20}$$

For $g = 0$ we get an ordinary differential equation that can be solved explicitly:

$$\widetilde{C}_0(s) = \frac{-1 + 18s + (1-12s)^{3/2}}{54s^2}\,.$$

It is easy to see that the substitution $s = t(1-3t)$ considerably simplifies $\widetilde{C}_0$ and makes it a rational function, namely

$$\widetilde{C}_0(t(1-3t)) = \frac{1-4t}{(1-3t)^2}$$

(cf. [10]). Substituting $s = t(1-3t)$ in (20), we get

$$\begin{aligned}
&t(1-2t)(1-3t)\,\dot{\widetilde{C}}_g + (1 - 4t + 6t^2)\widetilde{C}_g = t^2(1-3t)^2\\
&\times\left(\widetilde{D}_t\widetilde{C}_{g-1} + 3\sum_{i=1}^{g-1}\left(\widetilde{C}_i + \frac{t(1-3t)}{1-6t}\dot{\widetilde{C}}_i\right)\left(\widetilde{C}_{g-i} + \frac{t(1-3t)}{1-6t}\dot{\widetilde{C}}_{g-i}\right)\right),
\end{aligned} \tag{21}$$

where $\dot{\widetilde{C}}_g = \frac{d}{dt}\widetilde{C}_g(t(1-3t))$ and

$$\widetilde{D}_t = \frac{4t^3(1-3t)^3}{(1-6t)^3}\cdot\frac{d^3}{dt^3} + \frac{24t^2(1-3t)^2(1-9t+27t^2)}{(1-6t)^4}\cdot\frac{d^2}{dt^2}$$

$$+\frac{9t(1-3t)(3-56t+456t^2-1728t^3+2592t^4)}{(1-6t)^5}\cdot\frac{d}{dt}+3\,. \quad (22)$$

Assuming that $\widetilde{C}_0,\dots,\widetilde{C}_{g-1}$ are known, we can think of (21) as an ODE for $\widetilde{C}_g$. The integrating factor for this equation is $\frac{t(1-2t)}{1-3t}$, so that we get from (21)

$$\widetilde{C}_g(t(1-3t)=\frac{1-3t}{t(1-2t)}$$
$$\times\int t^2\left(\widetilde{D}_t\widetilde{C}_{g-1}+3\sum_{i=1}^{g-1}\left(\widetilde{C}_i+\frac{t(1-3t)}{1-6t}\dot{\widetilde{C}}_i\right)\left(\widetilde{C}_{g-i}+\frac{t(1-3t)}{1-6t}\dot{\widetilde{C}}_{g-i}\right)\right)dt\,. \quad (23)$$

Since by definition $\widetilde{C}_g(0)=0$ for all $g\geq 1$, Eq. (23) determines $\widetilde{C}_g$ uniquely in terms of $\widetilde{C}_0,\dots,\widetilde{C}_{g-1}$. In particular, this equation immediately yields

$$\widetilde{C}_1(t(1-3t))=\frac{t^2}{(1-2t)(1-6t)^2}\,.$$

Let us show that $\widetilde{C}_g(t(1-3t))$ has the form (17) for any $g\geq 2$. Using an analogue of (10) we obtain, after some cancellations, that the integrand in (23) has the form

$$\frac{\widetilde{Q}_g(t)}{(1-2t)^{3g-2}(1-6t)^{5g-2}}\,, \quad (24)$$

where $\widetilde{Q}_g(t)=\sum_{i=2g}^{8g-6}\tilde{q}_{g,i}t^i$ is a polynomial with $\tilde{q}_{g,2g}=(2g-1)(4g-1)(4g-3)\,p_{g-1,2g-2}$. It can be further decomposed into the sum

$$\frac{\widetilde{Q}_g(t)}{(1-2t)^{3g-2}(1-6t)^{5g-2}}=\sum_{i=2}^{3g-2}\frac{\tilde{a}_i}{(1-2t)^i}+\sum_{j=2}^{5g-2}\frac{\tilde{b}_j}{(1-6t)^j}\,. \quad (25)$$

Integrating it, we obtain

$$\int\frac{\widetilde{Q}_g(t)}{(1-2t)^{3g-2}(1-6t)^{5g-2}}dt=\tilde{b}+\sum_{i=1}^{3g-3}\frac{\tilde{a}_{i+1}}{2i}\cdot\frac{1}{(1-2t)^i}+\sum_{j=2}^{5g-3}\frac{\tilde{b}_{j+1}}{6j}\cdot\frac{1}{(1-6t)^j}\,, \quad (26)$$

where the condition $\widetilde{C}_g(0)=0$ implies

$$\tilde{b}=-\sum_{i=1}^{3g-3}\frac{\tilde{a}_{i+1}}{2i}-\sum_{j=1}^{5g-3}\frac{\tilde{b}_{j+1}}{6j}\,.$$

Multiplying the right hand side of (26) by $(1-2t)^{3g-3}(1-6t)^{5g-3}$ we get a polynomial of the form $\widetilde{R}_g(t)=\sum_{i=2g+1}^{8g-6}\tilde{r}_{g,i}t^i$. To complete the proof we put $\widetilde{P}_g(t)=\frac{1-3t}{t}\widetilde{R}_g(t)$ and notice that $\tilde{p}_{g,2g}=\frac{(2g-1)(4g-1)(4g-3)}{2g+1}\,\tilde{p}_{g-1,2g-2}$. Moreover, we see that $t=1/3$ is a root of $\widetilde{P}_g(t)$.[5]

□

Acknowledgements. This work was supported by the Russian Science Foundation: Theorem 1 by the grant 14-21-00035, and Theorem 2 by the grant 16-11-10316. We are grateful to F. Petrov for a useful suggestion. We thank A. Giorgetti, A. Mednykh, R. Nedela and T. R. S. Walsh for discussions.

[5]Numerically, we also have $\tilde{p}_{g,8g-6}\neq 0,\ \widetilde{P}_g(1/2)\neq 0,\ \widetilde{P}_g(1/6)\neq 0$.

References

[1] Adrianov, N.: An analog of the Harer–Zagier formula for unicellular bicolored maps, Func. Anal. Appl. **31**:3, 1–9 (1997).

[2] Carrell, S.R., Chapuy, G.: Simple recurrence formulas to count maps on orientable surfaces. J. Combinatorial Theory, Series A, **133**, 58–75 (2015).

[3] Giorgetti, A., Walsh, T. R. S.: Enumeration of hypermaps of a given genus, arXiv:1510.09019 (2015).

[4] Harer, J., Zagier, D.: The Euler characteristic of the moduli space of curves. Invent. Math. **85**:3, 457–485 (1986).

[5] Jackson, D.M.: Counting cycles in permutations by group characters, with an application to a topological problem. Trans. Amer. Math. Soc. **299**, 785–801 (1987).

[6] Kazarian, M., Zograf, P.: Virasoro constraints and topological recursion for Grothendiecks dessin counting. Lett. Math. Phys. **105** (8), 1057-1084 (2015).

[7] Lando, S. K., Zvonkin, A. K.: Graphs on surfaces and their applications. Encyclopaedia of Mathematical Sciences **141**, Springer-Verlag, Berlin (2004).

[8] Mednykh, A., Nedela, R.: Enumeration of unrooted maps of a given genus. J. Combinatorial Theory B **96**:5, 706–729 (2006).

[9] Mednykh, A., Nedela, R.: Enumeration of unrooted hypermaps of a given genus. Discrete Mathematics **310**:3, 518–526 (2010).

[10] Tutte, W. T.: A census of planar maps. Canadian Math. J. **15**:2, 249-271 (1963).

[11] Walsh, T. R. S.: Hypermaps versus bipartite maps. J. Combinatorial Theory B **18**:2, 155–163 (1975).

[12] Walsh, T. R. S., Lehman, A. B.: Counting rooted maps by genus. I. J. Combinatorial Theory B **13**, 192–218 (1972).

[13] Zograf, P.: Enumeration of Grothendieck's dessins and KP hierarchy, Int. Math. Res. Notices **2015**, 13533-13544 (2015).

Steklov Mathematical Institute, 8 Gubkin St., Moscow 119991 Russia, and Department of Mathematics, National Research University Higher School of Economics, Usacheva Str. 6, Moscow 119048 Russia

E-mail address: kazarian@mccme.ru

St.Petersburg Department of the Steklov Mathematical Institute, Fontanka 27, St. Petersburg 191023 Russia, and Chebyshev Laboratory of St. Petersburg State University, 14th Line V.O. 29B, St.Petersburg 199178 Russia

E-mail address: zograf@pdmi.ras.ru